

# **Coastal Wetland Habitat Dynamics in Selected New South Wales Estuaries.**

## **Volume 1**

Submitted by Kylee Margaret Wilton  
Bachelor of Arts  
Master of Science

A thesis submitted in total fulfilment of the requirements of  
the degree of Doctor of Philosophy.

School of Arts and Sciences (NSW)  
Faculty of Arts and Sciences

Australian Catholic University  
Research Services  
Locked Bay 4115  
Fitzroy, Victoria 3065

February, 2002

## Abstract

---

Intertidal wetland habitats in southeastern Australia have changed significantly during the past sixty years. Mangrove habitats have expanded both seawards and landwards, the latter being at the expense of saltmarsh habitats. This relatively common phenomenon is generally suggested to be an outcome of sea-level rise. Several factors potentially responsible for this change are examined, including changes in mean sea-level during the past 50 to 100 years, changes in climate, population growth, catchment landuse, and estuary type.

A protocol for mapping estuarine habitats was developed and implemented, incorporating the application of geographic information systems. Spatial and temporal coastal wetland habitat changes at nine sites along the New South Wales coast are illustrated. These habitat dynamics were shown to not correlate between sites. The results demonstrate that sea-level rise in this region cannot solely account for the extent of change during the past sixty years. With the exception of one site (Careel Bay), there have been no correlations between contemporary mean sea-level rise and mangrove incursion of the saltmarsh habitats at the study sites, or with rainfall patterns, at the scale of observation in this study, which was largely decadal. The only correlations determined during this study have been between population growth and coastal wetland habitat dynamics in some sites.

In spite of saltmarsh habitat loss being a regional phenomenon, local factors appear to have a profound bearing on the rates of change. Neither contemporary mean sea-level rise, rainfall patterns, estuary type, catchment landuse, catchment natural cover nor population pressure can account solely for the patterns in the spatial and temporal dynamics of the coastal wetlands of New South Wales. It seems apparent that regional factors create preconditions favourable for mangrove incursion, but that localised conditions have been responsible for the extent of these incursions from site to site. That is, despite higher sea-level and greater rainfall, the extent of change has been determined by the unique characteristics of each site.

The results have important implications for current estuary management practices in the state of New South Wales. The lack of spatial and temporal trends in coastal wetland habitat dynamics point to the need for management to be conducted on a localised, rather than regional scale. Additionally, anthropogenic influences must be carefully managed, since the extent of mangrove habitat expansion into saltmarsh areas is unlikely to be a natural occurrence.

This thesis contains no material extracted in whole or in part from a thesis by which I have qualified for or been awarded another degree or diploma. Chapter 3: Habitat Mapping Protocols, written specifically for this thesis, has been reproduced in its original form as an Australian Catholic University Technical Report.

No other person's work has been used without due acknowledgment in the main text of the thesis.

This thesis has not been submitted for the award of any degree or diploma in any other tertiary institution.

.....

Kylee M. Wilton

February 2002

date

## Acknowledgements

---

As for many Ph.D. candidates, this has been a long road, along which I have grown in many ways, some of which were unexpected. It has been an interesting experience that has helped me to adapt to many different situations and ways of thinking.

I would firstly like to acknowledge the support of the Australian Research Council and the Department of Employment, Education, Training and Youth Affairs, who provided the scholarship for this research, and to thank the Australian Catholic University for facilitating it. This scholarship was enhanced by the wonderful intellectual, practical and financial support of the Estuaries Branch of the New South Wales Department of Land and Water Conservation. I am indebted in many ways to this group, and hope to return to it sometime in the future. They made me feel a part of the team.

I would like to thank Dr Neil Saintilan of the Australian Catholic University, my principal supervisor, for giving me the wonderful opportunity of being a Ph.D. candidate, which has enabled me to explore a part of this magnificent coastline of New South Wales. His comments particularly during the draft phase of the project were most helpful. I would like to thank Dr Peter Cowell of the University of Sydney for firstly inspiring me as an undergraduate student like no other lecturer did, and for providing support when needed during this research. He taught me to approach science in new ways. I will forever be indebted to my third supervisor, Mr Bruce Coates of the Estuaries Branch of the New South Wales Department of Land and Water Conservation, who has provided me with an enormous amount of support, in many different ways. He believed in me, and supported me when I did not think I could make it to the end. I could not respect a colleague more. Thank you Bruce.

I am indebted to Kerrylee Rogers, of the Coastal Wetlands Unit of the Australian Catholic University, for seeing me through perhaps the hardest parts of this research. She helped me in all of my field work, and we spent many

hours wandering the mangroves and saltmarshes of New South Wales contemplating life, and getting very muddy. She also assisted me enormously in my attempts to master the Geographic Information System *ArcView*.

Many people helped me in the technical aspects of the research. Mr John Floyd of the Estuaries Branch of the Department of Land and Water Conservation answered my cries for computer technical support. Dr David Rissik of the same group helped me work my way through the thesis-writing process, and reviewed some drafts for me. I am also thankful to Dr Pia Laegdsgaard of DLWC, my fellow saltmarsh enthusiast, for keeping me sane at all the conferences and meetings.

I have received much-needed support from a number of other people indirectly related to my research, and I am truly thankful to them for their input to my endeavours.

On a personal note, I am thankful to all the people beyond the scope of this scientific pursuit who showed an interest in my work. It certainly helps, when you spend the majority of the time mapping (or designing golf courses, as the guys in the Estuaries Branch would say), or traipsing around in knee-deep mud in wetsuit booties all day! I am also terribly thankful to my small group of close friends who know and understand me, and who have provided incredible support in simple ways when times were tough.

My parents, Neville and Patricia Potts, have been a wonderful support, and have been since the beginning. I'm sure they didn't expect me to be 27 years old before I finished school! I love my parents deeply, they are my heroes, and I will forever be indebted to them for providing me with every possible opportunity. I cannot say enough about them.

Finally, my wonderful, wonderful husband Ross has endured this journey with me, and felt both my pain and enthusiasm. He is looking forward to getting his wife back! At any rate, he is my best friend, and has made a point of listening to me talk about my work, of understanding it, and providing objective feedback.

He has been my field assistant at times, but more often than not he has been there to provide me with logic during those times when it disappeared. He knows I am not the best with words, but I will be forever indebted to him for giving me all the support I needed to make it through this research. I know he, like my parents, wants me to make the most of every opportunity I receive, and he is the man behind the woman!

This research has felt like an enormous battle at times, but I know I am a better person for having made it through this far!

## TABLE OF CONTENTS

---

Abstract.....	i
Declaration.....	iii
Acknowledgements.....	iv
Table of Contents.....	vii
List of Figures.....	xi
List of Tables.....	xvii

### **CHAPTER 1: INTRODUCTION AND LITERATURE REVIEW..... 1**

1.1	Introduction	
1.1.1	Model of the response of saline coastal wetlands to changes in mean sea-level in New South Wales estuaries.....	1
1.1.2	Mangrove and saltmarsh distribution and ecology.....	4
1.1.2.1	Mangrove Habitats.....	5
1.1.2.2	Saltmarsh Habitats.....	9
1.1.2.3	The Mangrove and Saltmarsh Habitat Boundary.....	10
1.1.3	The Nature of Change in Intertidal Habitats.....	11
1.1.3.1	Conceptual Models of Intertidal Habitat Response to Sea-level Change.....	12
1.1.3.2	Stratigraphic Palaeo-scale Models of Intertidal Habitat Change.....	16
1.1.3.3	Contemporary Studies of Intertidal Habitat Change in Response to Sedimentation and Sea-level Change.....	17
1.1.4	Mangrove Transgression in Australasia.....	23
1.1.4.1	Trends Throughout Australia.....	25
1.1.4.2	Trends in New South Wales.....	26
1.1.4.3	Previous Studies of Wetland Habitat Change in Eastern Australia.....	27
1.1.4.4	Northern New South Wales.....	30
1.1.4.5	Central New South Wales.....	31
1.1.4.6	The Sydney Metropolitan Area.....	35
1.1.4.7	Southern New South Wales.....	38
1.1.4.8	Southern Australia.....	40
1.1.4.9	New Zealand.....	41
1.2	Aims and Hypotheses.....	43
1.3	Significance of This Study.....	44
1.4	Thesis Structure.....	45

### **CHAPTER 2: SITE SELECTION..... 47**

2.1	Criteria for Site Selection.....	47
2.1.1	New South Wales Estuaries.....	48
2.1.2	Rainfall.....	56
2.1.3	Population Change.....	61
2.2	Site Selection and Justification.....	69
2.3	Site Descriptions.....	71
2.3.1	Ukerebagh Island, Tweed River.....	71
2.3.2	Tilligerry Creek, Port Stephens.....	76

2.3.3	Black Neds Bay, Swansea Channel, Lake Macquarie.....	80
2.3.4	Pelican, Rileys and St Huberts Islands, Brisbane Water.....	85
2.3.5	Courangra Point, Hawkesbury River.....	88
2.3.6	Careel Bay, Pittwater.....	93
2.3.7	Towra Point, Botany Bay.....	98
2.3.8	Currambene Creek and Carama Inlet, Jervis Bay.....	102
2.4	Summary.....	107

### **CHAPTER 3: WETLAND HABITAT MAPPING PROTOCOLS..... 108**

3.1	Introduction.....	108
3.1.1	Objectives.....	108
3.1.2	Inventory vs. Monitoring Mapping.....	109
3.2	Previous Exercised in Mangrove and Saltmarsh Mapping.....	110
3.3	Methods.....	111
3.3.1	Data Input.....	111
3.3.1.1	Data Sources.....	111
3.3.1.2	Scale Rectification.....	111
3.3.1.2.1	Variation in Scale.....	111
3.3.1.2.2	Correcting Scale Using GIS Programs.....	112
3.3.1.2.3	Correcting Scale Using the AC3 Stereo Plotter.....	114
3.3.1.3	Community Classification.....	116
3.3.1.3.1	The Mangrove-Saltmarsh Boundary.....	116
3.3.1.3.2	The Saltmarsh-Terrestrial Boundary.....	118
3.3.1.3.3	Interpretation of Aerial Photographs.....	118
3.3.1.3.4	Supervised Classification.....	119
3.3.1.4	Community Delineation.....	119
3.3.1.4.1	Hand-drawn Boundaries.....	125
3.3.1.4.2	On-screen Digitising.....	125
3.3.1.5	Examples of On-screen Digitising.....	126
3.3.2	Data Analysis.....	129
3.3.2.1	Non-digital Analyses.....	129
3.3.2.2	Raster and Vector GIS Analyses.....	130
3.3.2.3	Cumulative Errors.....	132
3.3.3	Comparison of Techniques.....	132
3.4	Recommendations.....	134
3.5	Discussion.....	135

### **CHAPTER 4: WETLAND HABITAT MAPPING RESULTS.....139**

4.1	Introduction.....	139
4.2	Aims.....	139
4.3	Methods.....	140
4.3.1	Mapping.....	140
4.3.2	Data Derived From Mapping Calculations.....	144
4.4	Results.....	146
4.4.1	Maps of Coastal Wetland Habitat Areas and Changes.....	147
4.4.2	Coastal Wetland Habitat Areal Changes.....	165
4.4.3	Modes of Habitat Changes in Coastal Wetlands.....	169

4.4.4	Inter-Site Comparisons of Coastal Wetland Habitat Changes.....	188
4.5	Discussion: Trends in Coastal Wetland Habitat Dynamics in Selected New South Wales Estuaries.....	197
4.5.1	Introduction.....	197
4.5.2	Site-Specific Coastal Wetland Habitat Dynamics.....	199
4.5.3	Comparisons of Trends in Coastal Wetland Habitat Dynamics.....	211

**CHAPTER 5: SITE CONTOUR MODELLING AND THE EFFECT OF  
SEA-LEVEL RISE.....215**

5.1	Sea Level Rise.....	215
5.1.1	The Nature of Sea Level Rise.....	215
5.1.2	Global Sea Level Patterns.....	217
5.1.3	Australian Sea Level Patterns.....	218
5.1.4	Regional (Australian East Coast) Sea Level Patterns.....	219
5.2	Site Contour Development and Statistical Analyses.....	223
5.2.1	Introduction.....	223
5.2.2	Methods.....	224
5.2.2.1	Site Contour Development.....	224
5.2.2.2	Elevational Comparisons.....	232
5.2.2.3	Statistical Methods.....	232
5.2.3	Digital Elevation Models.....	236
5.2.4	Results of Site Contour Development.....	239
5.2.5	Results of Statistical Analyses of Coastal Wetland Habitat Elevations.....	242
5.2.5.1	Habitat Inundation Depths.....	242
5.2.5.2	Mean Sea-level Versus Mangrove Incursion.....	244
5.3	Discussion of Trends in Coastal Wetland Habitat Elevations.....	248

**CHAPTER 6: ASSESSMENT OF OTHER CONSIDERATIONS.....255**

6.1	Other Considerations.....	255
6.2	Climate Change.....	255
6.2.1	Introduction.....	255
6.2.2	Definitions.....	256
6.2.2.1	Air Temperature.....	256
6.2.2.2	El Niño.....	257
6.2.2.3	The Southern Oscillation Index.....	258
6.2.2.4	ENSO.....	259
6.2.2.5	Climatic Influences on Wetlands.....	260
6.2.2.6	Rainfall.....	260
6.2.2.7	Objectives.....	262
6.2.3	Rainfall and Mean Sea-level.....	263
6.2.3.1	Introduction.....	263
6.2.3.2	Aim.....	263
6.2.3.3	Methods.....	263
6.2.3.4	Results.....	264
6.2.3.5	Discussion.....	266
6.2.4	Rainfall and Habitat Change.....	267
6.2.4.1	Introduction.....	267

6.2.4.2	Aim.....	268
6.2.4.3	Methods.....	268
6.2.4.4	Results.....	270
6.2.4.5	Discussion.....	272
6.3	Catchment Landuse and Anthropogenic Influence.....	274
6.3.1	The Characteristics of Urbanisation.....	274
6.3.1.1	Introduction.....	274
6.3.1.2	Aim.....	278
6.3.1.3	Methods.....	278
6.3.1.4	Results.....	280
6.3.1.5	Discussion.....	282
<b>CHAPTER 7: DISCUSSION AND CONCLUSIONS.....</b>		<b>283</b>
7.1	Introduction.....	283
7.2	Review of Major Findings.....	283
7.2.1	Themes.....	283
7.2.1.1	Review of Aims.....	287
7.2.1.2	Review of Hypotheses.....	287
7.2.2	Coastal Wetland Habitat Dynamics.....	291
7.3	Concluding Remarks and Implications for Research and Management.....	300
7.3.1	Implications for Research.....	301
7.3.2	Implications for Management.....	302
<b>REFERENCES.....</b>		<b>305</b>
<b>APPENDIX 1: CATCHMENT TYPES.....</b>		<b>330</b>
<b>APPENDIX 2: AERIAL PHOTOGRAPH INDEX.....</b>		<b>332</b>
<b>APPENDIX 3: ARCVIEW MAPPING INSTRUCTIONS.....</b>		<b>338</b>
<b>APPENDIX 4: AERIAL PHOTOGRAPH GEORECTIFICATION ERROR.....</b>		<b>358</b>
<b>APPENDIX 5: COMPLETE STUDY SITE MAPS.....</b>		<b>360</b>
<b>APPENDIX 6: HABITAT MAPPING RAW DATASETS.....</b>		<b>450</b>
<b>APPENDIX 7: SITE CONTOUR MODELLING AND SEA LEVEL RAW DATASETS AND STATISTICS.....</b>		<b>497</b>
<b>APPENDIX 8: RAW DATASETS FOR ANALYSIS OF OTHER CONSIDERATIONS.....</b>		<b>522</b>
<b>APPENDIX 9: PREVIOUS PUBLICATIONS.....</b>		<b>528</b>
<b>APPENDIX 10: JOURNAL PAPER – SAINTILAN AND WILTON (2001)....</b>		<b>530</b>

## List of Figures

---

Figure No.	Figure Title	Page No.
1.1	Cross section of a typical wetland.....	5
1.2	Relationships between mangrove floristic, structural and functional diversity and the range of factors chiefly influencing mangrove distribution globally, regionally, locally and intertidally.....	9
2.1	NSW IBRA bioregions.....	53
2.2	NSW IMCRA bioregions.....	55
2.3	Average annual rainfall at Tweed Heads (Tweed Heads Golf Club), 1887 to 1997.....	57
2.4	Average annual rainfall in the Tilligerry Creek region (Williamstown), 1944 to 1998.....	57
2.5	Average annual rainfall in the Black Neds Bay area (Newcastle), 1862 to 1998.....	58
2.6	Average annual rainfall in the Brisbane Water area (Gosford), 1916 to 1998.....	58
2.7	Average annual rainfall in the Courangra Point area (Wisemans Ferry), 1903 to 1998.....	59
2.8	Average annual rainfall in Pittwater. Rainfall History for Newport (1932 to 1992) and Avalon (1994 to 1996).....	59
2.9	Average annual rainfall in the Towra Point area (Botany Bay, Sydney Airport), 1930 to 1998.....	60
2.10	Average annual rainfall in Jervis Bay, 1899 to 1998.....	60
2.11	Rates of population change, coastal statistical divisions, 1971 to 1998.....	62
2.12	Population levels in the Tweed Shire Council local government area, 1947 to 1999.....	62
2.13	Population levels in the Port Stephens local government area, 1947 to 1999.....	63
2.14	Population levels in the Lake Macquarie local government area, 1947 to 1999.....	63
2.15	Population levels in the Gosford local government area, 1947 to 1999.....	64
2.16	Population levels in the Pittwater area, 1947 to 1999.....	64
2.17	Population levels in the Botany (1947 to 1999) and Sutherland (1981 to 1999) local government areas .....	65
2.18	Population levels in the Shoalhaven local government area (1947 to 1998).....	65
2.19	The hydrologic cycle in a natural catchment.....	67
2.20	The hydrologic cycle in an anthropogenically-influenced catchment..	67
2.21	Study site selection along the New South Wales coast.....	69
2.22	Ukerebagh Island in the mouth of the Tweed River.....	71
2.23	Tilligerry Creek, in Port Stephens.....	77
2.24a	Black Neds Bay, in Swansea Channel, the mouth of Lake Macquarie.....	81
2.24b	Black Neds Bay.....	81
2.25	The study islands in Brisbane Water.....	86

2.26	Courangra Point, on the Hawkesbury River.....	89
2.27	Careel Bay, Pittwater.....	94
2.28	Towra Point, Botany Bay.....	98
2.29	Currambene Creek and Carama Inlet, Jervis Bay.....	104
3.1	Jervis Bay (1993) showing A: saltmarsh; B: mixed mangrove and saltmarsh; C: mangrove; D: <i>Casuarina</i> .....	120
3.2	Zone A: Saltmarsh ( <i>Sarcocornia quinqueflora</i> ) in the foreground, with individual mature mangrove trees in the background.....	121
3.3	<i>Sarcocornia quinqueflora</i> .....	121
3.4	Zone A: Saltmarsh ( <i>Juncus kraussii</i> ) in the foreground.....	122
3.5	<i>Juncus kraussii</i> .....	122
3.6	Zone B: Mixed mangrove and saltmarsh habitat.....	123
3.7	Zone B: Another example of mixed mangrove and saltmarsh habitat.....	123
3.8	Zone C: Mangrove woodland ( <i>Avicennia marina</i> ).....	124
3.9	Zone D: <i>Casuarina</i> woodland.....	124
3.10a	Currambene Creek, Jervis Bay, 1949.....	128
3.10b	Digitising of selected habitats in Currambene Creek, Jervis Bay, 1949.....	128
3.11a	Currambene Creek, Jervis Bay, 1993.....	129
3.11b	Digitising of selected habitats in Currambene Creek, Jervis Bay, 1993.....	129
3.12	Change in area (ha) of mangroves in Berowra and Marramarra Creeks, 1941-1994.....	133
4.1	Wetland habitats of Ukerebagh Island, Tweed River, 1948.....	147
4.2	Wetland habitats of Ukerebagh Island, Tweed River, 1998.....	147
4.3	Wetland habitat change on Ukerebagh Island, Tweed River, from 1948 to 1998.....	148
4.4	Gross wetland habitat change on Ukerebagh Island, Tweed Heads, 1948 to 1998.....	148
4.5	Wetland habitats at the mouth of Tilligerry Creek, Port Stephens, 1954.....	149
4.6	Wetland habitats at the mouth of Tilligerry Creek, Port Stephens, 1996.....	149
4.7	Wetland habitat change at the mouth of Tilligerry Creek, Port Stephens, 1954 to 1996.....	150
4.8	Gross wetland habitat change at the mouth of Tilligerry Creek, Port Stephens, 1954 to 1996.....	150
4.9	Wetland habitats in Black Neds Bay, Swansea Channel, Lake Macquarie, 1950.....	151
4.10	Wetland habitats in Black Neds Bay, Swansea Channel, Lake Macquarie, 1999.....	151
4.11	Wetland habitat change in Black Neds Bay, Swansea Channel, Lake Macquarie, 1950 to 1999.....	152
4.12	Gross wetland habitat change in Black Neds Bay, Swansea Channel, Lake Macquarie, 1950 to 1999.....	152
4.13	Selected wetland habitats in Brisbane Water, 1954.....	153
4.14	Selected wetland habitats in Brisbane Water, 1998.....	153

4.15	Wetland habitat change on selected islands in Brisbane Water, 1954 to 1998.....	154
4.16	Gross wetland habitat change at selected sites in Brisbane Water, 1954 to 1998.....	154
4.17	Wetland habitats at Courangra Point, Hawkesbury River, 1954.....	155
4.18	Wetland habitats at Courangra Point, Hawkesbury River, 1994.....	155
4.19	Wetland habitat change at Courangra Point, Hawkesbury River, 1954 to 1994.....	156
4.20	Gross wetland habitat change at Courangra Point, Hawkesbury River, 1954 to 1994.....	156
4.21	Wetland habitats in Careel Bay, Pittwater, 1940.....	157
4.22	Wetland habitats in Careel Bay, Pittwater, 1996.....	157
4.23	Wetland habitat change in Careel Bay, Pittwater, 1940 to 1996.....	158
4.24	Gross wetland habitat change in Careel Bay, Pittwater, 1940 to 1996.....	158
4.25	Wetland habitats at Towra Point, Botany Bay, 1956.....	159
4.26	Wetland habitats at Towra Point, Botany Bay, 1999.....	159
4.27	Habitat change at Towra Point, Botany Bay, 1956 to 1999.....	160
4.28	Gross wetland habitat change at Towra Point, Botany Bay, 1956 to 1999.....	160
4.29	Wetland habitats in Currumbene Creek, Jervis Bay, 1949.....	161
4.30	Wetland habitats in Currumbene Creek, Jervis Bay, 1993.....	161
4.31	Wetland habitat change in Currumbene Creek, Jervis Bay, 1949 to 1993.....	162
4.32	Gross wetland habitat change in Currumbene Creek, Jervis Bay, 1949 to 1993.....	162
4.33	Wetland habitats in Cararma Inlet, Jervis Bay, 1948.....	163
4.34	Wetland habitats in Cararma Inlet, Jervis Bay, 1999.....	163
4.35	Wetland habitat change in Cararma Inlet, Jervis Bay, 1948 to 1999.....	164
4.36	Gross wetland habitat change in Cararma Inlet, Jervis Bay, 1948 to 1999.....	164
4.37	Absolute values of the main modes of habitat change on Ukerebagh Island, Tweed River, 1948 to 1998.....	170
4.38a	The fate of the original saltmarsh habitat on Ukerebagh Island from 1948 to 1998, based on the original 13.3 ha in 1948.....	171
4.38b	Habitat types lost at the expense of areas of new mangrove habitat (11.1 ha) on Ukerebagh Island, from 1948 to 1998.....	171
4.39	Absolute values of the main modes of habitat change in the mouth of Tilligerry Creek, Port Stephens, 1954 to 1996.....	172
4.40a	The fate of the original saltmarsh habitat in the mouth of Tilligerry Creek from 1954 to 1996, based on the original 222.5 ha in 1954...	173
4.40b	Habitat types lost at the expense of areas of new mangrove habitat (83.7 ha) in the mouth of Tilligerry Creek from 1954 to 1996.....	173
4.41	Absolute values of the main modes of habitat change in Black Neds Bay, Lake Macquarie, 1950 to 1999.....	174
4.42a	The fate of the original saltmarsh habitat in Black Neds Bay from 1950 to 1999, based on the original 38.5 ha in 1950.....	175
4.42b	Habitat types lost at the expense of areas of new mangrove habitat (39.8 ha) in Black Neds Bay from 1950 to 1999.....	175

4.43	Absolute values of the main modes of habitat change on selected islands in Brisbane Water, 1954 to 1998.....	176
4.44a	The fate of the original saltmarsh habitat on selected islands in Brisbane Water from 1954 to 1998, based on the original 44.3 ha in 1954.....	177
4.44b	Habitat types lost at the expense of areas of new mangrove habitat (31.4 ha) on selected islands in Brisbane Water from 1954 to 1998..	177
4.45	Absolute values of the main modes of habitat change at Courangra Point, Hawkesbury River, 1954 to 1994.....	178
4.46a	The fate of the original saltmarsh habitat at Courangra Point, Hawkesbury River from 1954 to 1994, based on the original 47.7 ha in 1954.....	179
4.46b	Habitat types lost at the expense of areas of new mangrove habitat (21.2 ha) at Courangra Point, Hawkesbury River from 1954 to 1994.....	179
4.47	Absolute values of the main modes of habitat change in Careel Bay, Pittwater, 1940 to 1996.....	180
4.48a	The fate of the original saltmarsh habitat in Careel Bay, Pittwater, 1940 to 1996, based on the original 21.6 ha in 1940.....	181
4.48b	Habitat types lost at the expense of areas of new mangrove habitat (9 ha) in Careel Bay, Pittwater, 1940 to 1996.....	181
4.49	Absolute values of the main modes of habitat change at Towra Point, Botany Bay, 1956 to 1999.....	182
4.50a	The fate of the original saltmarsh habitat at Towra Point, Botany Bay from 1956 to 1999, based on the original 229.3 ha in 1956.....	183
4.50b	Habitat types lost at the expense of areas of new mangrove habitat (152.6 ha) at Towra Point, Botany Bay, 1956 to 1999.....	183
4.51	Absolute values of the main modes of habitat change in Currumbene Creek, Jervis Bay, 1949 to 1993.....	184
4.52a	The fate of the original saltmarsh habitat in Currumbene Creek, Jervis Bay from 1949 to 1993, based on the original 51 ha in 1949..	185
4.52b	Habitat types lost at the expense of areas of new mangrove habitat (30.4 ha) in Currumbene Creek, Jervis Bay, 1949 to 1993.....	185
4.53	Absolute values of the main modes of habitat change in Cararma Inlet, Jervis Bay, 1948 to 1999.....	186
4.54a	The fate of the original saltmarsh habitat in Cararma Inlet, Jervis Bay from 1948 to 1999, based on the original 181.7 ha in 1948.....	187
4.54b	Habitat types lost at the expense of areas of new mangrove habitat (44.5 ha) in Cararma Inlet, Jervis Bay, 1948 to 1999.....	187
4.55a	Multidimensional scaling of all changes, as a percentage of all changes within each site, to show relationships between sites.....	190
4.55b	Multidimensional scaling of all changes, as a percentage of all changes within each site, with an explanation of relationships between sites.....	191
4.56	Multidimensional scaling, showing the relative importance of the seaward expansion of mangrove habitat into the adjacent waterway.....	192
4.57	Multidimensional scaling, showing the relative importance of change from mixed to mangrove habitat.....	193

4.58	Multidimensional scaling, showing the relative presence of change from saltmarsh to mangrove habitat.....	194
4.59	Multidimensional scaling, showing the relative presence of change from saltmarsh to mixed habitat.....	195
4.60	Multidimensional scaling, showing the relative presence of terrestrial incursion into saltmarsh habitat.....	196
4.61	Multidimensional scaling, showing the relative presence of saltmarsh incursion into terrestrial habitat.....	197
5.1	Mean sea-level in Sydney (Fort Denison), 1915 to 2000.....	220
5.2	Tidal characteristics of a drowned river valley estuary (Hawkesbury River).....	222
5.3	Tidal characteristics of a tidal river (Manning River).....	222
5.4	Tidal characteristics of a tidal lake (Lake Macquarie).....	222
5.5	Accuracy of tidal inundation modelling at Towra Point over four consecutive high tides.....	225
5.6	Example of dowel and transect placement within a study site.....	226
5.7	Illustration of how measurements are taken to produce a digital elevation model of the site.....	226
5.8	Location of dowel transects in Black Neds Bay.....	226
5.9	Extrapolated data points in Black Neds Bay based on original dowel transects.....	226
5.10	Location of dowel transects at Courangra Point.....	227
5.11	Extrapolated data at Courangra Point based on original dowel transects.....	227
5.12	Location of dowel transects in Careel Bay.....	227
5.13	Extrapolated data points in Careel Bay based on original dowel transects.....	227
5.14	Location of dowel transects in Towra Point.....	228
5.15	Close-up of location of dowel transects in Towra Point.....	228
5.16	Extrapolated data points in Towra Point based on original dowel transects.....	228
5.17	Location of dowel transects in Currambene Creek.....	229
5.18	Extrapolated data points in Currambene Creek based on original dowel transects.....	229
5.19	A digital elevation model produced (this example is Currambene Creek, Jervis Bay) after inserting field dowel elevation information into the GIS <i>ArcView</i> .....	230
5.20	Model of the changes in an average wetland profile, 1940s/50s to 1990s.....	232
5.21	Digital elevation model of Black Neds Bay.....	236
5.22	Digital elevation model of Courangra Point.....	237
5.23	Digital elevation model of Careel Bay.....	237
5.24	Digital elevation model of Towra Point.....	238
5.25	Digital elevation model of Currambene Creek.....	238
5.26	Mean depth below Spring high water of habitat maintenance and change in Black Neds Bay, Swansea Channel, Lake Macquarie, based on habitat change between 1950 and 1999.....	240

5.27	Mean depth below Spring high water of habitat maintenance and change at Courangra Point, Hawkesbury River, based on habitat change between 1954 and 1994.....	240
5.28	Mean depth below Spring high water of habitat maintenance and change in Careel Bay, Pittwater, based on habitat change between 1940 and 1996.....	241
5.29	Mean depth below Spring high water of habitat maintenance and change at Towra Point, Botany Bay, based on habitat change between 1956 and 1999.....	241
5.30	Mean depth below Spring high water of habitat maintenance and change in Currumbene Creek, Jervis Bay, based on habitat change between 1949 and 1993.....	242
5.31	The correlation between mean sea-level and the depth below Spring high water (2 m) of mangrove incursion, at all sites.....	244
5.32	The correlation between mean sea-level and the depth below Spring high water (2 m) of mangrove incursion 5 years later, at all sites.....	245
5.33	The correlation between mean sea-level and the mean depth below Spring high water (2 m) of the change in elevation of mangrove incursion, at all sites .....	245
5.34	The correlation between mean sea-level and the depth below Spring high water of the change in the elevation of mangrove incursion with a 5 year lag, at all sites.....	246
5.35	The correlation between mean sea-level and the depth below Spring high water (2 m) of mangrove incursion, at Careel Bay.....	247
5.36	The correlation between mean sea-level and the depth below Spring high water (2 m) of mangrove incursion, at Careel Bay, with a 5 year lag.....	247
6.1	The correlation between mean sea-level (Fort Denison) and rainfall in Tilligerry Creek, 1915 to 1997.....	265
6.2	The correlation between mean sea-level (Fort Denison) and rainfall in Black Neds Bay, 1915 to 1997.....	265
6.3	The correlation between mean sea-level (Fort Denison) and rainfall in Brisbane Water, 1915 to 1997.....	265
6.4	The correlation between mean sea-level (Fort Denison) and rainfall at Courangra Point, 1915 to 1997.....	265
6.5	The correlation between mean sea-level (Fort Denison) and rainfall in Careel Bay, 1915 to 1997.....	266
6.6	The correlation between mean sea-level (Fort Denison) and rainfall at Towra Point, 1915 to 1997.....	266
6.7	The correlation between mean sea-level (Fort Denison) and rainfall in Jervis Bay, 1915 to 1997.....	266
6.8	Rates of population changes, coastal statistical divisions, 1971 to 1998.....	275
6.9	Absolute change in population, coastal areas, 1994 to 1999.....	275
6.10	Population growth at the study sites.....	279

## List of Tables

---

Table No.	Table Title	Page No.
1.1	Studies reporting spatial and/or temporal changes in saltmarsh and mangrove habitats around the coast of Australia.....	23
1.2	Estimated areas (ha) of saltmarsh/saltpan habitat altered in Australian States.....	25
1.3	Estimated areas (ha) of mangrove habitat altered in Australian States .....	26
1.4	Recent wetland vegetation assessments in Queensland, which highlight several factors associated with change.....	30
2.1	Roy <i>et al.</i> 's (2001) grouping of estuary types in eastern Australia....	49
2.2	Roy <i>et al.</i> 's (2001) modified estuary classification scheme.....	50
2.3	Digby <i>et al.</i> 's (1999) criteria used for the classification of estuaries.....	51
2.4	Coastal wetland habitats chosen for habitat dynamics assessment....	52
2.5	Site selection for this study.....	70
3.1	Characteristics of raster and vector data structures used in geographic information systems.....	130
3.2	Change in area (ha) of mangroves in Berowra and Marramarra Creeks, 1941-1994.....	133
4.1	The area (ha) and percentage change (%) of individual habitat types on Ukerebagh Island from 1948 to 1998.....	165
4.2	The area (ha) and percentage change (%) of individual habitat types at the mouth of Tilligerry Creek from 1954 to 1996.....	165
4.3	The area (ha) and percentage change (%) of individual habitat types in Black Neds Bay from 1950 to 1999.....	166
4.4	The area (ha) and percentage change (%) of individual habitat types at selected sites in Brisbane Water from 1954 to 1998.....	166
4.5	The area (ha) and percentage change (%) of individual habitat types at Courangra Point from 1954 to 1994.....	167
4.6	The area (ha) and percentage change (%) of individual habitat types in Careel Bay from 1940 to 1996.....	167
4.7	The area (ha) and percentage change (%) of individual habitat types at Towra Point from 1956 to 1999.....	168
4.8	The area (ha) and percentage change (%) of individual habitat types in Currumbene Creek from 1949 to 1993.....	168
4.9	The area (ha) and percentage change (%) of individual habitat types in Cararma Inlet from 1948 to 1999.....	169
4.10	The area (ha) and percentage change (% , from period to period) of the primary modes of habitat change from period to period, 1948 to 1998, on Ukerebagh Island, Tweed River.....	170
4.11	The area (ha) and percentage change (% , from period to period) of the primary modes of habitat change from period to period, 1954 to 1996, in the mouth of Tilligerry Creek, Port Stephen.....	172

4.12	The area (ha) and percentage change (% , from period to period) of the primary modes of habitat change from period to period, 1950 to 1999, in Black Neds Bay, Lake Macquarie.....	174
4.13	The area (ha) and percentage change (% , from period to period) of the primary modes of habitat change from period to period, 1954 to 1998, on selected islands in Brisbane Water.....	176
4.14	The area (ha) and percentage change (% , from period to period) of the primary modes of habitat change from period to period, 1954 to 1994, at Courangra Point.....	178
4.15	The area (ha) and percentage change (% , from period to period) of the primary modes of habitat change from period to period, 1940 to 1996, in Careel Bay, Pittwater.....	180
4.16	The area (ha) and percentage change (% , from period to period) of the primary modes of habitat change from period to period, 1956 to 1999, at Towra Point, Botany Bay.....	182
4.17	The area (ha) and percentage change (% , from period to period) of the primary modes of habitat change from period to period, 1949 to 1993, in Currumbene Creek, Jervis Bay.....	184
4.18	The area (ha) and percentage change (% , from period to period) of the primary modes of habitat change from period to period, 1948 to 1999, in Cararma Inlet, Jervis Bay.....	186
4.19	The area (ha) and percentage increase (+) or decrease (-) (% , from period to period) of mangrove habitat at each site during each date/data set.....	188
4.20	The area (ha) and percentage increase (+) or decrease (-) (% , from period to period) of mixed habitat at each site during each date/data set. Note: net change is in the context of gains and losses in the mangrove and saltmarsh habitats, concurrently.....	189
4.21	The area (ha) and percentage increase (+) or decrease (-) (% , from period to period) of saltmarsh habitat at each site during each date/data set.....	189
5.1	Raw data for the correlation between the mean inundation depth of mangrove incursion below a 2 metre Spring high water, per site, per decade of observation.....	234
5.2	Raw data for the correlation between the mean change in inundation depth of mangrove incursion below a 2 metre Spring high water, per site, per decade of observation.....	235
5.3	Mean inundation depths of habitat zones (m) below Spring high water (2 m).....	239
5.4	Tests of Between Subject Effects based on elevation.....	243
5.5	Statistical results of the analysis of the mean depth below Spring high water (2 m) of vegetation zones at each site .....	244
5.6	The correlation between mean sea-level and the elevation of mangrove incursion at each site.....	248
5.7	Comparison of the elevational ranges of coastal wetland habitats, Clarke and Hannon (1969) versus this thesis.....	249

5.8	Comparison of the mean degree of mangrove incursion (i.e. the difference in depth of inundation between maintained mangrove habitat and mangrove incursion into saltmarsh habitat) and the increase in mean sea-level at each study site.....	254
6.1	Bureau of Meteorology data gauging site information.....	264
6.2	Site-specific correlation and regression data for the relationship between rainfall and mean sea-level, 1915 to 1997.....	265
6.3	Correlations between (a) mean annual rainfall for the 10 years previous to the mapping date, and corresponding mangrove habitat areas, and (b) 10 year mean annual rainfall and a 5-year lag in mangrove habitat areas.....	270
6.4	Correlations between (a) mean annual rainfall for the 10 years previous to the mapping date, and corresponding saltmarsh habitat areas, and (b) 10 year mean annual rainfall and a 5-year lag in saltmarsh habitat areas.....	270
6.5	Correlations between (a) mean annual rainfall and corresponding mangrove habitat gain, and (b) mean annual rainfall and a 5-year lag in mangrove habitat gain.....	271
6.6	Correlations between (a) mean annual rainfall and corresponding saltmarsh habitat loss, and (b) mean annual rainfall and a 5-year lag in saltmarsh habitat loss.....	271
6.7	Correlations between (a) mean annual rainfall and saltmarsh loss to mangrove incursion (%), and (b) mean annual rainfall and a 5-year lag in saltmarsh loss to mangrove incursion (%). ....	272
6.8	Correlations between (a) population total and corresponding mangrove area, and (b) population total and a 5-year lag in mangrove area.....	280
6.9	Correlations between (a) population total and corresponding saltmarsh area, and (b) population total and a 5-year lag in saltmarsh area.....	281
6.10	Correlations between (a) population growth and corresponding saltmarsh habitat loss to mangrove incursion (%), and (b) population growth and a 5-year lag in saltmarsh habitat loss to mangrove incursion (%). ....	281
7.1	Changes in the extent of saltmarsh habitats in Australasian estuaries.....	286
7.2	Comparison of wetland study site area, from smallest to largest.....	292
7.3	Comparison of key wetland habitat changes and factors.....	293
7.4	Comparison of percentage decline of absolute saltmarsh area, from least to greatest.....	292
7.5	Comparison of that proportion (%) of original saltmarsh habitat lost to mangrove incursion (gross), from least to greatest.....	294
7.6	Comparison of that proportion (%) of original saltmarsh habitat lost to reclamation (gross), from least to greatest.....	295
7.7	Comparison of that proportion (%) of original saltmarsh habitat lost to terrestrial incursion (gross), from least to greatest.....	295

7.8	Comparison of both the degree of catchment development (indicated by a tick mark in the appropriate column reflecting the proportion of catchment natural cover) and the degree of saltmarsh loss to mangrove incursion (% value shown), north to south.....	296
7.9	Comparison of both estuary type (indicated by a tick mark in the appropriate column) and the degree of saltmarsh loss to mangrove incursion (% value shown).....	297